

Package ‘SetTest’

July 21, 2025

Type Package

Title Group Testing Procedures for Signal Detection and Goodness-of-Fit

Version 0.3.0

Author Hong Zhang and Zheyang Wu

Maintainer Hong Zhang <hzhang@wpi.edu>

Description It provides cumulative distribution function (CDF), quantile, p-value, statistical power calculator and random number generator for a collection of group-testing procedures, including the Higher Criticism tests, the one-sided Kolmogorov-Smirnov tests, the one-sided Berk-Jones tests, the one-sided phi-divergence tests, etc. The input are a group of p-values. The null hypothesis is that they are i.i.d. Uniform(0,1). In the context of signal detection, the null hypothesis means no signals. In the context of the goodness-of-fit testing, which contrasts a group of i.i.d. random variables to a given continuous distribution, the input p-values can be obtained by the CDF transformation. The null hypothesis means that these random variables follow the given distribution. For reference, see [1]Hong Zhang, Jiashun Jin and Zheyang Wu. ``Distributions and power of optimal signal-detection statistics in finite case'', IEEE Transactions on Signal Processing (2020) 68, 1021-1033; [2] Hong Zhang and Zheyang Wu. ``The general goodness-of-fit tests for correlated data'', Computational Statistics & Data Analysis (2022) 167, 107379.

License GPL-2

Encoding UTF-8

RoxygenNote 7.2.0

NeedsCompilation no

Repository CRAN

Date/Publication 2024-07-15 03:40:02 UTC

Contents

pbj 2

phc	3
power.bj	4
power.hc	6
power.phi	7
pphi	9
pphi.omni	10
qbj	11
qhc	12
qphi	13
stat.bj	14
stat.hc	15
stat.phi	16
stat.phi.omni	17
test.bj	19
test.hc	20
test.phi	21
test.phi.omni	22

Index **24**

pbj *CDF of Berk-Jones statistic under the null hypothesis.*

Description

CDF of Berk-Jones statistic under the null hypothesis.

Usage

pbj(q, M, k0, k1, onesided = FALSE, method = "ecc", ei = NULL)

Arguments

q	- quantile, must be a scalar.
M	- correlation matrix of input statistics (of the input p-values).
k0	- search range starts from the k0th smallest p-value.
k1	- search range ends at the k1th smallest p-value.
onesided	- TRUE if the input p-values are one-sided.
method	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
ei	- the eigenvalues of M if available.

Value

The left-tail probability of the null distribution of B-J statistic at the given quantile.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

See Also

[stat.bj](#) for the definition of the statistic.

Examples

```
pval <- runif(10)
bjstat <- stat.phi(pval, s=1, k0=1, k1=10)$value
pbj(q=bjstat, M=diag(10), k0=1, k1=10)
```

phc

CDF of Higher Criticism statistic under the null hypothesis.

Description

CDF of Higher Criticism statistic under the null hypothesis.

Usage

```
phc(q, M, k0, k1, onesided = FALSE, method = "ecc", ei = NULL)
```

Arguments

- | | |
|----------|--|
| q | - quantile, must be a scalar. |
| M | - correlation matrix of input statistics (of the input p-values). |
| k0 | - search range starts from the k0th smallest p-value. |
| k1 | - search range ends at the k1th smallest p-value. |
| onesided | - TRUE if the input p-values are one-sided. |
| method | - default = "ecc": the effective correlation coefficient method in reference 2.
"ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method. |
| ei | - the eigenvalues of M if available. |

Value

The left-tail probability of the null distribution of HC statistic at the given quantile.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

See Also

[stat.hc](#) for the definition of the statistic.

Examples

```
pval <- runif(10)
hcstat <- stat.phi(pval, s=2, k0=1, k1=5)$value
phc(q=hcstat, M=diag(10), k0=1, k1=10)
```

power.bj

Statistical power of Berk and Jones test.

Description

Statistical power of Berk and Jones test.

Usage

```
power.bj(
  alpha,
  n,
  beta,
  method = "gaussian-gaussian",
  eps = 0,
  mu = 0,
  df = 1,
  delta = 0
)
```

Arguments

- | | |
|--------|--|
| alpha | - type-I error rate. |
| n | - dimension parameter, i.e. the number of input statistics to construct B-J statistic. |
| beta | - search range parameter. Search range = (1, beta*n). Beta must be between 1/n and 1. |
| method | - different alternative hypothesis, including mixtures such as, "gaussian-gaussian", "gaussian-t", "t-t", "chisq-chisq", and "exp-chisq". By default, we use Gaussian mixture. |

eps	- mixing parameter of the mixture.
mu	- mean of non standard Gaussian model.
df	- degree of freedom of t/Chisq distribution and exp distribution.
delta	- non-cenrality of t/Chisq distribution.

Details

We consider the following hypothesis test,

$$H_0 : X_i \sim F, H_a : X_i \sim G$$

Specifically, $F = F_0$ and $G = (1 - \epsilon)F_0 + \epsilon F_1$, where ϵ is the mixing parameter, F_0 and F_1 is speified by the "method" argument:

"gaussian-gaussian": F_0 is the standard normal CDF and $F = F_1$ is the CDF of normal distribution with μ defined by mu and $\sigma = 1$.

"gaussian-t": F_0 is the standard normal CDF and $F = F_1$ is the CDF of t distribution with degree of freedom defined by df.

"t-t": F_0 is the CDF of t distribution with degree of freedom defined by df and $F = F_1$ is the CDF of non-central t distribution with degree of freedom defined by df and non-centrality defined by delta.

"chisq-chisq": F_0 is the CDF of Chisquare distribution with degree of freedom defined by df and $F = F_1$ is the CDF of non-central Chisquare distribution with degree of freedom defined by df and non-centrality defined by delta.

"exp-chisq": F_0 is the CDF of exponential distribution with parameter defined by df and $F = F_1$ is the CDF of non-central Chisqaure distribution with degree of freedom defined by df and non-centrality defined by delta.

Value

Power of BJ test.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and Statistical Power of Optimal Signal-Detection Methods In Finite Cases", submitted.
2. Donoho, David; Jin, Jiashun. "Higher criticism for detecting sparse heterogeneous mixtures". Annals of Statistics 32 (2004).
3. Jager, Leah; Wellner, Jon A. "Goodness-of-fit tests via phi-divergences". Annals of Statistics 35 (2007).
4. Berk, R.H. & Jones, D.H. Z. "Goodness-of-fit test statistics that dominate the Kolmogorov statistics". Wahrscheinlichkeitstheorie verw Gebiete (1979) 47: 47.

See Also

[stat.bj](#) for the definition of the statistic.

Examples

```
power.bj(0.05, n=10, beta=0.5, eps = 0.1, mu = 1.2)
```

```
power.hc
```

Statistical power of Higher Criticism test.

Description

Statistical power of Higher Criticism test.

Usage

```
power.hc(
  alpha,
  n,
  beta,
  method = "gaussian-gaussian",
  eps = 0,
  mu = 0,
  df = 1,
  delta = 0
)
```

Arguments

alpha	- type-I error rate.
n	- dimension parameter, i.e. the number of input statistics to construct Higher Criticism statistic.
beta	- search range parameter. Search range = (1, beta*n). Beta must be between 1/n and 1.
method	- different alternative hypothesis, including mixtures such as, "gaussian-gaussian", "gaussian-t", "t-t", "chisq-chisq", and "exp-chisq". By default, we use Gaussian mixture.
eps	- mixing parameter of the mixture.
mu	- mean of non standard Gaussian model.
df	- degree of freedom of t/Chisq distribution and exp distribution.
delta	- non-cenrality of t/Chisq distribution.

Details

We consider the following hypothesis test,

$$H_0 : X_i \sim F, H_a : X_i \sim G$$

Specifically, $F = F_0$ and $G = (1 - \epsilon)F_0 + \epsilon F_1$, where ϵ is the mixing parameter, F_0 and F_1 is speified by the "method" argument:

"gaussian-gaussian": F_0 is the standard normal CDF and $F = F_1$ is the CDF of normal distribution with μ defined by mu and $\sigma = 1$.

"gaussian-t": F_0 is the standard normal CDF and $F = F_1$ is the CDF of t distribution with degree of freedom defined by df.

"t-t": F_0 is the CDF of t distribution with degree of freedom defined by df and $F = F_1$ is the CDF of non-central t distribution with degree of freedom defined by df and non-centrality defined by delta. "chisq-chisq": F_0 is the CDF of Chisquare distribution with degree of freedom defined by df and $F = F_1$ is the CDF of non-central Chisquare distribution with degree of freedom defined by df and non-centrality defined by delta.

"exp-chisq": F_0 is the CDF of exponential distribution with parameter defined by df and $F = F_1$ is the CDF of non-central Chisquare distribution with degree of freedom defined by df and non-centrality defined by delta.

Value

Power of HC test.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and Statistical Power of Optimal Signal-Detection Methods In Finite Cases", submitted.
2. Donoho, David; Jin, Jiashun. "Higher criticism for detecting sparse heterogeneous mixtures". Annals of Statistics 32 (2004).

See Also

[stat.hc](#) for the definition of the statistic.

Examples

```
power.hc(0.05, n=10, beta=0.5, eps = 0.1, mu = 1.2)
```

power.phi

Statistical power of phi-divergence test.

Description

Statistical power of phi-divergence test.

Usage

```
power.phi(
  alpha,
  n,
  s,
  beta,
  method = "gaussian-gaussian",
```

```

    eps = 0,
    mu = 0,
    df = 1,
    delta = 0
)

```

Arguments

alpha	- type-I error rate.
n	- dimension parameter, i.e. the number of input statistics to construct phi-divergence statistic.
s	- phi-divergence parameter. $s = 2$ is the higher criticism statistic. $s = 1$ is the Berk and Jones statistic.
beta	- search range parameter. Search range = $(1, \text{beta} \cdot n)$. Beta must be between $1/n$ and 1.
method	- different alternative hypothesis, including mixtures such as, "gaussian-gaussian", "gaussian-t", "t-t", "chisq-chisq", and "exp-chisq". By default, we use Gaussian mixture.
eps	- mixing parameter of the mixture.
mu	- mean of non standard Gaussian model.
df	- degree of freedom of t/Chisq distribution and exp distribution.
delta	- non-cenrality of t/Chisq distribution.

Details

We consider the following hypothesis test,

$$H_0 : X_i \sim F, H_a : X_i \sim G$$

Specifically, $F = F_0$ and $G = (1 - \epsilon)F_0 + \epsilon F_1$, where ϵ is the mixing parameter, F_0 and F_1 is speified by the "method" argument:

"gaussian-gaussian": F_0 is the standard normal CDF and $F = F_1$ is the CDF of normal distribution with μ defined by mu and $\sigma = 1$.

"gaussian-t": F_0 is the standard normal CDF and $F = F_1$ is the CDF of t distribution with degree of freedom defined by df.

"t-t": F_0 is the CDF of t distribution with degree of freedom defined by df and $F = F_1$ is the CDF of non-central t distribution with degree of freedom defined by df and non-centrality defined by delta.

"chisq-chisq": F_0 is the CDF of Chisquare distribution with degree of freedom defined by df and $F = F_1$ is the CDF of non-central Chisquare distribution with degree of freedom defined by df and non-centrality defined by delta.

"exp-chisq": F_0 is the CDF of exponential distribution with parameter defined by df and $F = F_1$ is the CDF of non-central Chisquare distribution with degree of freedom defined by df and non-centrality defined by delta.

Value

Power of phi-divergence test.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and Statistical Power of Optimal Signal-Detection Methods In Finite Cases", submitted.
2. Donoho, David; Jin, Jiashun. "Higher criticism for detecting sparse heterogeneous mixtures". Annals of Statistics 32 (2004).

See Also

[stat.phi](#) for the definition of the statistic.

Examples

```
#If the alternative hypothesis Gaussian mixture with eps = 0.1 and mu = 1.2:#
power.phi(0.05, n=10, s=2, beta=0.5, eps = 0.1, mu = 1.2)
```

pphi	<i>calculate the left-tail probability of phi-divergence under general correlation matrix.</i>
------	--

Description

calculate the left-tail probability of phi-divergence under general correlation matrix.

Usage

```
pphi(q, M, k0, k1, s = 2, t = 30, onesided = FALSE, method = "ecc", ei = NULL)
```

Arguments

- | | |
|----------|---|
| q | - quantile, must be a scalar. |
| M | - correlation matrix of input statistics (of the input p-values). |
| k0 | - search range starts from the k0th smallest p-value. |
| k1 | - search range ends at the k1th smallest p-value. |
| s | - the phi-divergence test parameter. |
| t | - numerical truncation parameter. |
| onesided | - TRUE if the input p-values are one-sided. |
| method | - default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method. |
| ei | - the eigenvalues of M if available. |

Value

Left-tail probability of the phi-divergence statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033 2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379 3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

Examples

```
M = toeplitz(1/(1:10)*(-1)^(0:9)) #alternating polynomial decaying correlation matrix
pphi(q=2, M=M, k0=1, k1=5, s=2)
pphi(q=2, M=M, k0=1, k1=5, s=2, method = "ecc")
pphi(q=2, M=M, k0=1, k1=5, s=2, method = "ave")
pphi(q=2, M=diag(10), k0=1, k1=5, s=2)
```

pphi.omni	<i>calculate the left-tail probability of omnibus phi-divergence statistics under general correlation matrix.</i>
-----------	---

Description

calculate the left-tail probability of omnibus phi-divergence statistics under general correlation matrix.

Usage

```
pphi.omni(q, M, K0, K1, S, t = 30, onesided = FALSE, method = "ecc", ei = NULL)
```

Arguments

q	- quantile, must be a scalar.
M	- correlation matrix of input statistics (of the input p-values).
K0	- vector of search range starts (from the k0th smallest p-value).
K1	- vector of search range ends (at the k1th smallest p-value).
S	- vector of the phi-divergence test parameters.
t	- numerical truncation parameter.
onesided	- TRUE if the input p-values are one-sided.
method	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
ei	- the eigenvalues of M if available.

Value

Left-tail probability of omnibus phi-divergence statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

Examples

```
M = matrix(0.3,10,10) + diag(1-0.3, 10)
pphi.omni(0.05, M=M, K0=rep(1,2), K1=rep(5,2), S=c(1,2))
```

qbj

Quantile of Berk-Jones statistic under the null hypothesis.

Description

Quantile of Berk-Jones statistic under the null hypothesis.

Usage

```
qbj(p, M, k0, k1, onesided = FALSE, method = "ecc", ei = NULL, err_thr = 1e-04)
```

Arguments

- | | |
|----------|---|
| p | - a scalar left probability that defines the quantile. |
| M | - correlation matrix of input statistics (of the input p-values). |
| k0 | - search range starts from the k0th smallest p-value. |
| k1 | - search range ends at the k1th smallest p-value. |
| onesided | - TRUE if the input p-values are one-sided. |
| method | - default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method. |
| ei | - the eigenvalues of M if available. |
| err_thr | - the error threshold. The default value is 1e-4. |

Value

Quantile of BJ statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

See Also

[stat.bj](#) for the definition of the statistic.

Examples

```
## The 0.05 critical value of BJ statistic when n = 10:
qbj(p=.95, M=diag(10), k0=1, k1=5, onesided=FALSE)
qbj(p=1-1e-5, M=diag(10), k0=1, k1=5, onesided=FALSE, err_thr=1e-8)
```

qhc

Quantile of Higher Criticism statistics under the null hypothesis.

Description

Quantile of Higher Criticism statistics under the null hypothesis.

Usage

```
qhc(p, M, k0, k1, onesided = FALSE, method = "ecc", ei = NULL, err_thr = 1e-04)
```

Arguments

- | | |
|----------|---|
| p | - a scalar left probability that defines the quantile. |
| M | - correlation matrix of input statistics (of the input p-values). |
| k0 | - search range starts from the k0th smallest p-value. |
| k1 | - search range ends at the k1th smallest p-value. |
| onesided | - TRUE if the input p-values are one-sided. |
| method | - default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method. |
| ei | - the eigenvalues of M if available. |
| err_thr | - the error threshold. The default value is 1e-4. |

Value

Quantile of HC statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

See Also

[stat.hc](#) for the definition of the statistic.

Examples

```
## The 0.05 critical value of HC statistic when n = 10:
qhc(p=.95, M=diag(10), k0=1, k1=5, onesided=FALSE)
qhc(p=1-1e-5, M=diag(10), k0=1, k1=5, onesided=FALSE, err_thr=1e-8)
```

qphi

Quantile of phi-divergence statistic under the null hypothesis.

Description

Quantile of phi-divergence statistic under the null hypothesis.

Usage

```
qphi(
  p,
  M,
  k0,
  k1,
  s = 2,
  t = 30,
  onesided = FALSE,
  method = "ecc",
  ei = NULL,
  err_thr = 1e-04
)
```

Arguments

- p - a scalar left probability that defines the quantile.
- M - correlation matrix of input statistics (of the input p-values).
- k0 - search range starts from the k0th smallest p-value.
- k1 - search range ends at the k1th smallest p-value.
- s - the phi-divergence test parameter.

t	- numerical truncation parameter.
onesided	- TRUE if the input p-values are one-sided.
method	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
ei	- the eigenvalues of M if available.
err_thr	- the error threshold. The default value is 1e-4.

Value

Quantile of the phi-divergence statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033 2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379 3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

See Also

[stat.phi](#) for the definition of the statistic.

Examples

```

qphi(p=.95, M=diag(10), k0=1, k1=5, s=2, onesided=FALSE)
qphi(p=1-1e-3, M=diag(10), k0=1, k1=5, s=2, onesided=FALSE)
qphi(p=1-1e-3, M=diag(10), k0=1, k1=5, s=2, onesided=FALSE, err_thr = 1e-6)
qphi(p=1-1e-5, M=diag(10), k0=1, k1=5, s=2, onesided=FALSE)
qphi(p=1-1e-5, M=diag(10), k0=1, k1=5, s=2, onesided=FALSE, err_thr = 1e-6)
qphi(p=1-1e-5, M=diag(10), k0=1, k1=5, s=2, onesided=FALSE, err_thr = 1e-8)

```

stat.bj

Construct Berk and Jones (BJ) statistics.

Description

Construct Berk and Jones (BJ) statistics.

Usage

```
stat.bj(p, k0 = 1, k1 = NA)
```

Arguments

- p - vector of input p-values.
k0 - search range left end parameter. Default k0 = 1.
k1 - search range right end parameter. Default k1 = 0.5*number of input p-values.

Details

Let $p_{(i)}$, $i = 1, \dots, n$ be a sequence of ordered p-values, the Berk and Jones statistic

$$BJ = \sqrt{2n} \max_{1 \leq i \leq \lfloor \beta n \rfloor} (-1)^j \sqrt{i/n * \log(i/n/p_{(i)}) + (1 - i/n) * \log((1 - i/n)/(1 - p_{(i)}))}$$

and when $p_{(i)} > i/n$, $j = 1$, otherwise $j = 0$.

Value

- value - BJ statistic constructed from a vector of p-values.
location - the order of the p-values to obtain BJ statistic.
stat - vector of marginal BJ statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and Statistical Power of Optimal Signal-Detection Methods In Finite Cases", submitted.
2. Jager, Leah; Wellner, Jon A. "Goodness-of-fit tests via phi-divergences". *Annals of Statistics* 35 (2007).
3. Berk, R.H. & Jones, D.H. Z. "Goodness-of-fit test statistics that dominate the Kolmogorov statistics". *Wahrscheinlichkeitstheorie verw Gebiete* (1979) 47: 47.

Examples

```
stat.bj(runif(10))
#When the input are statistics#
stat.test = rnorm(20)
p.test = 1 - pnorm(stat.test)
stat.bj(p.test, k0 = 2, k1 = 20)
```

stat.hc

Construct Higher Criticism (HC) statistics.

Description

Construct Higher Criticism (HC) statistics.

Usage

```
stat.hc(p, k0 = 1, k1 = NA)
```

Arguments

- p - vector of input p-values.
k0 - search range left end parameter. Default k0 = 1.
k1 - search range right end parameter. Default k1 = 0.5*number of input p-values.

Details

Let $p_{(i)}$, $i = 1, \dots, n$ be a sequence of ordered p-values, the higher criticism statistic

$$HC = \sqrt{n} \max_{1 \leq i \leq \lfloor \beta n \rfloor} [i/n - p_{(i)}] / \sqrt{p_{(i)}(1 - p_{(i)})}$$

Value

- value - HC statistic constructed from a vector of p-values.
location - the order of the p-values to obtain HC statistic.
stat - vector of marginal HC statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and Statistical Power of Optimal Signal-Detection Methods In Finite Cases", submitted.
2. Donoho, David; Jin, Jiashun. "Higher criticism for detecting sparse heterogeneous mixtures". Annals of Statistics 32 (2004).

Examples

```
stat.hc(runif(10))
#When the input are statistics#
stat.test = rnorm(20)
p.test = 1 - pnorm(stat.test)
stat.hc(p.test, k0 = 1, k1 = 10)
```

stat.phi

Construct phi-divergence statistics.

Description

Construct phi-divergence statistics.

Usage

```
stat.phi(p, s, k0 = 1, k1 = NA)
```


Arguments

- p - vector of input p-values.
- s - phi-divergence parameter. s = 2 is the higher criticism statistic. s = 1 is the Berk and Jones statistic.
- k0 - search range left end parameter. Default k0 = 1.
- k1 - search range right end parameter. Default k1 = 0.5*number of input p-values.

Details

Let $p_{(i)}$, $i = 1, \dots, n$ be a sequence of ordered p-values, the phi-divergence statistic

$$PHI = \sqrt{2n/(s - s^2)} \max_{1 \leq i \leq \lfloor \beta n \rfloor} (-1)^j \sqrt{1 - (i/n)^s (p_{(i)})^s - (1 - i/n)^{(1-s)} * (1 - p_{(i)})^{(1-s)}}$$

and when $p_{(i)} > i/n$, $j = 1$, otherwise $j = 0$.

Value

value - phi-divergence statistic constructed from a vector of p-values.

location - the order of the p-values to obtain phi-divergence statistic.

stat - vector of marginal phi-divergence statistics.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and Statistical Power of Optimal Signal-Detection Methods In Finite Cases", submitted.
2. Jager, Leah; Wellner, Jon A. "Goodness-of-fit tests via phi-divergences". Annals of Statistics 35 (2007).

Examples

```
stat.phi(runif(10), s = 2)
#When the input are statistics#
stat.test = rnorm(20)
p.test = 1 - pnorm(stat.test)
stat.phi(p.test, s = 0.5, k0 = 2, k1 = 5)
```

stat.phi.omni	<i>calculate the omnibus phi-divergence statistics under general correlation matrix.</i>
---------------	--

Description

calculate the omnibus phi-divergence statistics under general correlation matrix.

Usage

```
stat.phi.omni(
  p,
  M,
  K0 = rep(1, 2),
  K1 = rep(length(M[1, ]), 2),
  S = c(1, 2),
  t = 30,
  onesided = FALSE,
  method = "ecc",
  ei = NULL
)
```

Arguments

p	- input pvalues.
M	- correlation matrix of input statistics (of the input p-values).
K0	- vector of search range starts (from the k0th smallest p-value).
K1	- vector of search range ends (at the k1th smallest p-value).
S	- vector of the phi-divergence test parameters.
t	- numerical truncation parameter.
onesided	- TRUE if the input p-values are one-sided.
method	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
ei	- the eigenvalues of M if available.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033 2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379 3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

Examples

```
p.test = runif(10)
M = toeplitz(1/(1:10)*(-1)^(0:9)) #alternating polynomial decaying correlation matrix
stat.phi.omni(p.test, M=M, K0=rep(1,2), K1=rep(5,2), S=c(1,2))
```

`test.bj`*Multiple comparison test using Berk and Jones (BJ) statistics.*

Description

Multiple comparison test using Berk and Jones (BJ) statistics.

Usage

```
test.bj(prob, M, k0, k1, onesided = FALSE, method = "ecc", ei = NULL)
```

Arguments

<code>prob</code>	- vector of input p-values.
<code>M</code>	- correlation matrix of input statistics (of the input p-values).
<code>k0</code>	- search range starts from the <code>k0</code> th smallest p-value.
<code>k1</code>	- search range ends at the <code>k1</code> th smallest p-value.
<code>onesided</code>	- TRUE if the input p-values are one-sided.
<code>method</code>	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
<code>ei</code>	- the eigenvalues of <code>M</code> if available.

Value

`pvalue` - the p-value of the Berk-Jones test.

`bjstat` - the Berk-Jones statistic.

`location` - the order of the input p-values to obtain BJ statistic.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.
4. Leah Jager and Jon Wellner. "Goodness-of-fit tests via phi-divergences". Annals of Statistics 35 (2007).

See Also

[stat.bj](#) for the definition of the statistic.

Examples

```
test.bj(runif(10), M=diag(10), k0=1, k1=10)
#When the input are statistics#
stat.test = rnorm(20)
p.test = 2*(1 - pnorm(abs(stat.test)))
test.bj(p.test, M=diag(20), k0=1, k1=10)
```

test.hc

*Multiple comparison test using Higher Criticism (HC) statistics.***Description**

Multiple comparison test using Higher Criticism (HC) statistics.

Usage

```
test.hc(prob, M, k0, k1, onesided = FALSE, method = "ecc", ei = NULL)
```

Arguments

prob	- vector of input p-values.
M	- correlation matrix of input statistics (of the input p-values).
k0	- search range starts from the k0th smallest p-value.
k1	- search range ends at the k1th smallest p-value.
onesided	- TRUE if the input p-values are one-sided.
method	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
ei	- the eigenvalues of M if available.

Value

pvalue - The p-value of the HC test.

hcstat - HC statistic.

location - the order of the input p-values to obtain HC statistic.

References

- Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
- Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
- Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.
- Donoho, David; Jin, Jiashun. "Higher criticism for detecting sparse heterogeneous mixtures". Annals of Statistics 32 (2004).

See Also

[stat.hc](#) for the definition of the statistic.

Examples

```
pval.test = runif(10)
test.hc(pval.test, M=diag(10), k0=1, k1=10)
#When the input are statistics#
stat.test = rnorm(20)
p.test = 2*(1 - pnorm(abs(stat.test)))
test.hc(p.test, M=diag(20), k0=1, k1=10)
```

test.phi

Multiple comparison test using phi-divergence statistics.

Description

Multiple comparison test using phi-divergence statistics.

Usage

```
test.phi(prob, M, k0, k1, s = 2, onesided = FALSE, method = "ecc", ei = NULL)
```

Arguments

prob	- vector of input p-values.
M	- correlation matrix of input statistics (of the input p-values).
k0	- search range starts from the k0th smallest p-value.
k1	- search range ends at the k1th smallest p-value.
s	- phi-divergence parameter. s = 2 is the higher criticism statistic. s = 1 is the Berk and Jones statistic.
onesided	- TRUE if the input p-values are one-sided.
method	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
ei	- the eigenvalues of M if available.

Value

pvalue - The p-value of the phi-divergence test.

phistat - phi-divergence statistic.

location - the order of the input p-values to obtain phi-divergence statistic.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.
4. Leah Jager and Jon Wellner. "Goodness-of-fit tests via phi-divergences". Annals of Statistics 35 (2007).

See Also

[stat.phi](#) for the definition of the statistic.v

Examples

```
stat.test = rnorm(20) # Z-scores
p.test = 2*(1 - pnorm(abs(stat.test)))
test.phi(p.test, M=diag(20), s = 0.5, k0=1, k1=10)
test.phi(p.test, M=diag(20), s = 1, k0=1, k1=10)
test.phi(p.test, M=diag(20), s = 2, k0=1, k1=10)
```

test.phi.omni	<i>calculate the right-tail probability of omnibus phi-divergence statistics under general correlation matrix.</i>
---------------	--

Description

calculate the right-tail probability of omnibus phi-divergence statistics under general correlation matrix.

Usage

```
test.phi.omni(prob, M, K0, K1, S, onesided = FALSE, method = "ecc", ei = NULL)
```

Arguments

prob	- vector of input p-values.
M	- correlation matrix of input statistics (of the input p-values).
K0	- vector of search range starts (from the k0th smallest p-value).
K1	- vector of search range ends (at the k1th smallest p-value).
S	- vector of the phi-divergence test parameters.
onesided	- TRUE if the input p-values are one-sided.
method	- default = "ecc": the effective correlation coefficient method in reference 2. "ave": the average method in reference 3, which is an earlier version of reference 2. The "ecc" method is more accurate and numerically stable than "ave" method.
ei	- the eigenvalues of M if available.

Value

p-value of the omnibus test.

p-values of the individual phi-divergence test.

References

1. Hong Zhang, Jiashun Jin and Zheyang Wu. "Distributions and power of optimal signal-detection statistics in finite case", IEEE Transactions on Signal Processing (2020) 68, 1021-1033
2. Hong Zhang and Zheyang Wu. "The general goodness-of-fit tests for correlated data", Computational Statistics & Data Analysis (2022) 167, 107379
3. Hong Zhang and Zheyang Wu. "Generalized Goodness-Of-Fit Tests for Correlated Data", arXiv:1806.03668.

Examples

```
M = matrix(0.3,10,10) + diag(1-0.3, 10)
test.phi.omni(runif(10), M=M, K0=rep(1,2), K1=rep(5,2), S=c(1,2))
```

Index

pbj, 2
phc, 3
power.bj, 4
power.hc, 6
power.phi, 7
pphi, 9
pphi.omni, 10

qbj, 11
qhc, 12
qphi, 13

stat.bj, 3, 5, 12, 14, 19
stat.hc, 4, 7, 13, 15, 21
stat.phi, 9, 14, 16, 22
stat.phi.omni, 17

test.bj, 19
test.hc, 20
test.phi, 21
test.phi.omni, 22