

Exact unconditional sampling from max-stable random vectors

Léo Belzile

EPFL-SB-MATH-STAT
Station 8, Bâtiment MA
CH-1015, Switzerland,
leo.belzile@epfl.ch

The ‘mev’ package was originally introduced to implement the exact unconditional sampling algorithms in Dombry et al. (2016). The two algorithms therein allow one to simulate simple max-stable random vectors. The implementation will work efficiently for moderate dimensions and do not currently take approximate schemes for large spatial domains.

1 Functions and use

There are two main functions, `rmev` and `rmevspec`. `rmev` samples from simple max-stable processes, meaning it will return an $n \times d$ matrix of samples, where each of the column has a sample from a unit Fréchet distribution. In contrast, `rmevspec` returns sample on the unit simplex from the spectral (or angular) measure. One could use this to test estimation based on spectral densities, or to construct samples from Pareto processes.

The syntax is

```
library(mev)
#Sample of size 1000 from a 5-dimensional logistic model
x <- rmev(n=1000, d=5, param=0.5, model="log")
#Marginal parameters are all standard Fréchet, meaning GEV(1,1,1)
apply(x, 2, function(col){ismev::gev.fit(col, show=FALSE)$mle})

##      [,1] [,2] [,3] [,4] [,5]
## [1,] 1.0240 1.008 1.0800 0.9986 1.011
## [2,] 1.0115 1.025 1.0626 1.0024 1.002
## [3,] 0.9758 1.023 0.9808 1.0051 1.004

#Sample from the corresponding spectral density
w <- rmevspec(n=1000, d=5, param=0.5, model="log")
#All rows sum to 1 by construction
head(rowSums(w))

## [1] 1 1 1 1 1
```

```
#The marginal mean is 1/d
round(colMeans(w),2)

## [1] 0.19 0.20 0.20 0.20 0.20
```

2 Description of the models implemented

The different models implemented are described in Dombry et al. (2016), but some other models can be found and are described here.

1. **logistic distribution** (log) The logistic model of Gumbel (1960) (the Gumbel Archimedean copula)

$$P(\mathbf{X} \leq \mathbf{x}) = \exp \left[- \left(\sum_{i=1}^n \left(\frac{1}{x_i} \right)^\alpha \right)^{\frac{1}{\alpha}} \right]$$

for $\alpha > 1$. By default, `rmev` will transform an argument in $(0, 1)$ without warning, to conform with the implementation. The spectral measure density is

$$h_{\mathbf{W}}(\mathbf{w}) = \frac{1}{d} \frac{\Gamma(d-\alpha)}{\Gamma(1-\alpha)} \alpha^{d-1} \left(\prod_{j=1}^d w_j \right)^{-(\alpha+1)} \left(\sum_{j=1}^d w_j^{-\alpha} \right)^{1/\alpha-d}, \quad \mathbf{w} \in \mathbb{S}_d$$

2. **asymmetric logistic distribution** (alog) This model was proposed by Tawn (1990). It shares the same parametrization as the `evd` package, merely replacing the algorithm for the generation of logistic variates. Let \mathbb{B}_d be the collection of all nonempty subsets of $\{1, \dots, d\}$. The distribution function of the d -variate asymmetric logistic distribution is

$$P(\mathbf{X} \leq \mathbf{x}) = \exp \left[- \sum_{b \in \mathbb{B}_d} \left(\sum_{i \in b} \left(\frac{\theta_{i,b}}{x_i} \right)^{\alpha_b} \right)^{\frac{1}{\alpha_b}} \right],$$

while the spectral density is

$$h_{\mathbf{W}}(\mathbf{w}) = \frac{1}{d} \frac{\prod_{j=1}^{d-1} (j\alpha_b - 1)}{\prod_{i \in b} w_i} \left(\prod_{i \in b} \frac{\theta_{i,b}}{w_i} \right)^{\alpha_b} \left(\sum_{i \in b} \left(\frac{\theta_{i,b}}{w_i} \right)^{\alpha_b} \right)^{1/\alpha_b-d}$$

The parameters $\theta_{i,b}$ must be provided in a list and represent the asymmetry parameter. The sampling algorithm, from Stephenson (2003) gives some insight on the construction mechanism as a max-mixture of logistic distributions. Consider sampling \mathbf{Z}_b from a logistic distribution of dimension $|b|$ (or Fréchet variates if $|b| = 1$) with parameter α_b (possibly recycled). Each marginal value corresponds to the maximum of the weighted corresponding entry. That is, $X_i = \max_{b \in \mathbb{B}_d} \theta_{i,b} Z_{i,b}$ for all $i = 1, \dots, d$. The max-mixture is valid provided that $\sum_{b \in \mathbb{B}_d} \theta_{i,b} = 1$ for $i = 1, \dots, d$. As such, empirical estimates of the spectral measure will almost surely place mass on the inside of the simplex rather than on subfaces.

3. **negative logistic distribution** (`neglog`) The distribution function of the min-stable distribution due to Galambos (1975) is

$$P(\mathbf{X} \leq \mathbf{x}) = \exp \left[- \sum_{b \in \mathbb{B}} (-1)^{|b|} \left(\sum_{i \in b} x_i^{-\alpha} \right)^{\frac{1}{\alpha}} \right]$$

for $\alpha \leq 0$ and corresponds to the model in Dombry et al. (2016). The density is (Kotz and Nadarajah, 2000)

$$h_{\mathbf{W}}(\mathbf{w}) = \frac{1}{d} \sum_{\substack{b \in \mathbb{B} \\ c \subset b}} (-1)^{|b|} \left(\prod_{j=1}^{d-1} (1 - j\alpha) \right) \left(\prod_{i \in c} w_j \right)^{-(\alpha+1)} \left(\sum_{i \in c} w_i^{-\alpha} \right)^{1/\alpha-d}$$

4. **asymmetric negative logistic distribution** (`aneglog`) The asymmetric negative logistic model is alluded to in Joe (1990) as a generalization of the Galambos model. It is constructed in the same way as the asymmetric logistic distribution; see Theorem 1 in Stephenson (2003). Let $\alpha_b \leq 0$ for all $b \in \mathbb{B}_d$ and $\theta_{i,b} \geq 0$ with $\sum_{b \in \mathbb{B}_d} \theta_{i,b} = 1$; the distribution function is then

$$P(\mathbf{X} \leq \mathbf{x}) = \exp \left[- \sum_{b \in \mathbb{B}} \sum_{c \in b} (-1)^{|c|} \left(\sum_{i \in c} \left(\frac{\theta_{i,b}}{x_i} \right)^{\alpha_b} \right)^{\frac{1}{\alpha_b}} \right].$$

The density is (Kotz and Nadarajah, 2000)

$$h_{\mathbf{W}}(\mathbf{w}) = \frac{1}{d} \sum_{\substack{b \in \mathbb{B} \\ c \subset b}} (-1)^{|b|} \left(\prod_{j=1}^{d-1} (1 - j\alpha) \right) \left(\prod_{i \in c} w_j \right)^{-(\alpha+1)} \left(\prod_{j \in b} \theta_{j,b} \right)^{\alpha_b} \left(\sum_{i \in c} \left(\frac{\theta_{i,b}}{w_i} \right)^{\alpha_b} \right)^{1/\alpha_b-d}$$

In particular, it does not correspond to the “negative logistic distribution” given in e.g. §4.2 of Coles and Tawn (1991) or §3.5.3 of Kotz and Nadarajah (2000). The latter is not a valid distribution function in dimension $d \geq 3$ as the constraints therein on the parameters $\theta_{i,b}$ are necessary, but not sufficient.

Joe (1990) mentions generalizations of the distribution as given above but the constraints were not enforced elsewhere in the literature. The proof that the distribution is valid follows from Theorem 1 of Stephenson (2003) as it is a max-mixture. Note that the parametrization of the asymmetric negative logistic distribution does not match the bivariate implementation of `rbvevd`.

5. **multilogistic distribution** (`bilog`) This multivariate extension of the logistic, proposed by Boldi (2009), places mass on the interior of the simplex. Let $\mathbf{W} \in \mathbb{S}_d$ be the solution of

$$\frac{W_j}{W_d} = \frac{C_j U_j^{-\alpha_j}}{C_d U_d^{-\alpha_d}}, \quad j = 1, \dots, d$$

where $C_j = \Gamma(d - \alpha_j) / \Gamma(1 - \alpha_j)$ for $j = 1, \dots, d$ and $\mathbf{U} \in \mathbb{S}_d$ follows a d -mixture of Dirichlet with the j th component being $\mathcal{D}(\mathbf{1} - \delta_j \alpha_j)$, so that the mixture has density function

$$h_{\mathbf{U}}(\mathbf{u}) = \frac{1}{d} \sum_{j=1}^d \frac{\Gamma(d - \alpha_j)}{\Gamma(1 - \alpha_j)} u_j^{-\alpha_j}$$

for $0 < \alpha_j < 1, j = 1, \dots, d$. The spectral density is given by

$$h_{\mathbf{W}}(\mathbf{w}) = \frac{1}{d} \left(\sum_{j=1}^d \alpha_j u_j \right)^{-1} \left(\prod_{j=1}^d \alpha_j u_d \right) \left(\sum_{j=1}^d \frac{\Gamma(d - \alpha_j)}{\Gamma(1 - \alpha_j)} u_j^{-\alpha_j} \right) \prod_{j=1}^d w_j^{-1}$$

for $\alpha_j \in (0, 1)$

6. **Coles and Tawn Dirichlet distribution** (ct) The Dirichlet model of Coles and Tawn (1991)

$$h_{\mathbf{W}}(\mathbf{w}) = \frac{1}{d} \frac{\Gamma\left(1 + \sum_{j=1}^d \alpha_j\right)}{\prod_{j=1}^d \alpha_j w_j} \left(\sum_{j=1}^d \alpha_j w_j \right)^{-(d+1)} \prod_{j=1}^d \alpha_j \prod_{j=1}^d \left(\frac{\alpha_j w_j}{\sum_{k=1}^d \alpha_k w_k} \right)^{\alpha_j - 1}$$

for $\alpha_j > 0$.

7. **scaled extremal Dirichlet** (sdir) The angular density of the scaled extremal Dirichlet model with parameters $\rho > -\min(\boldsymbol{\alpha})$ and $\boldsymbol{\alpha} \in \mathbb{R}_+^d$ is given, for all $\mathbf{w} \in \mathcal{S}_d$, by

$$h_{\mathbf{W}}(\mathbf{w}) = \frac{\Gamma(\bar{\alpha} + \rho)}{d \rho^{d-1} \prod_{i=1}^d \Gamma(\alpha_i)} \langle \{c(\boldsymbol{\alpha}, \rho)\}^{1/\rho}, \mathbf{w}^{1/\rho} \rangle^{-\rho - \bar{\alpha}} \prod_{i=1}^d \{c(\alpha_i, \rho)\}^{\alpha_i / \rho} w_i^{\alpha_i / \rho - 1}.$$

where $c(\boldsymbol{\alpha}, \rho)$ is the d -vector with entries $\Gamma(\alpha_i + \rho) / \Gamma(\alpha_i)$ for $i = 1, \dots, d$.

8. **Hüsler–Reiss** (hr), due to Hüsler and Reiss (1989). It is a special case of the Brown–Resnick process. While Engelke et al. (2015) state that Hüsler–Reiss variates can be sampled following the same scheme, the spatial analog is conditioned on a particular site (\mathbf{s}_0), which complicates the comparisons with the other methods.

Let $I_{-j} = \{1, \dots, d\} \setminus \{j\}$ and $\lambda_{ij}^2 \geq 0$ be entries of a strictly conditionally negative definite matrix $\boldsymbol{\Lambda}$, for which $\lambda_{ij}^2 = \lambda_{ji}^2$. Then, following Nikoloulopoulos et al. (2009) (Remark 2.5) and Huser and Davison (2013), we can write the distribution function as

$$\mathbb{P}(\mathbf{X} \leq \mathbf{x}) = \exp \left[- \sum_{j=1}^d \frac{1}{x_j} \Phi_{d-1, \boldsymbol{\Sigma}_{-j}} \left(\lambda_{ij} - \frac{1}{2\lambda_{ij}} \log \left(\frac{x_j}{x_i} \right), i \in I_{-j} \right) \right].$$

where the partial correlation matrix $\boldsymbol{\Sigma}_{-j}$ has elements

$$\varrho_{i,k;j} = \frac{\lambda_{ij}^2 + \lambda_{kj}^2 - \lambda_{ik}^2}{2\lambda_{ij}\lambda_{kj}}$$

and $\lambda_{ii} = 0$ for all $i \in I_{-j}$ so that the diagonal entries $\varrho_{i,i;j} = 1$.¹

The `evd` package implementation has a bivariate implementation of the Hüsler–Reiss distribution with dependence parameter r , with $r_{ik} = 1/\lambda_{ik}$ or $2/r = \sqrt{2\gamma(\mathbf{h})}$ for $\mathbf{h} = \|\mathbf{s}_i - \mathbf{s}_k\|$ for the

¹Engelke et al. (2015) uses the covariance matrix with entries are $\varsigma = 2(\lambda_{ij}^2 + \lambda_{kj}^2 - \lambda_{ik}^2)$, so the resulting expression is evaluated at $2\lambda_{ij}^2 - \log\left(\frac{x_j}{x_i}\right)$ instead. We recover the same expression by standardizing, since this amounts to division by the standard deviations $2\lambda_{ij}$.

Brown–Resnick model. In this setting, it is particularly easy since the only requirement is non-negativity of the parameter. For inference in dimension $d > 2$, one needs to impose the constraint $\Lambda = \{\lambda_{ij}^2\}_{i,j=1}^d \in \mathcal{D}$ (cf. Engelke et al. (2015), p.3), where

$$\mathcal{D} = \left\{ \mathbf{A} \in [0, \infty)^{d \times d} : \mathbf{x}^\top \mathbf{A} \mathbf{x} < 0, \forall \mathbf{x} \in \mathbb{R}^d \setminus \{\mathbf{0}\} \right. \\ \left. \text{with } \sum_{i=1}^d x_i = 0, a_{ij} = a_{ji}, a_{ii} = 0 \forall i, j \in \{1, \dots, d\} \right\}$$

denotes the set of symmetric conditionally negative definite matrices with zero diagonal entries. An avenue to automatically satisfy these requirements is to optimize over a symmetric positive definite matrix parameter $\Sigma = \mathbf{L}^\top \mathbf{L}$, where \mathbf{L} is an upper triangular matrix whose diagonal elements are on the log-scale to ensure uniqueness of the Cholesky factorization; see Pinheiro and Bates (1996). By taking

$$\Lambda(\Sigma) = \begin{pmatrix} 0 & \text{diag}(\Sigma)^\top \\ \text{diag}(\Sigma) & \mathbf{1} \text{diag}(\Sigma)^\top + \text{diag}(\Sigma) \mathbf{1}^\top - 2\Sigma \end{pmatrix}$$

one can perform unconstrained optimization for the non-zero elements of \mathbf{L} which are in one-to-one correspondence with those of Λ .

It easily follows that generating \mathbf{Z} from a $d - 1$ dimensional log-Gaussian distribution with covariance $\text{Cov}(Z_i, Z_k) = 2(\lambda_{ij}^2 + \lambda_{kj}^2 - \lambda_{ik}^2)$ for $i, k \in I_{-j}$ with mean vector $-2\lambda_{\cdot j}^2$ gives the finite dimensional analog of the Brown–Resnick process in the mixture representation of Dombry et al. (2016).

The `rmev` function checks conditional negative definiteness of the matrix. The easiest way to do so negative definiteness of Λ with real entries is to form $\tilde{\Lambda} = \mathbf{P}\Lambda\mathbf{P}^\top$, where \mathbf{P} is an $d \times d$ matrix with ones on the diagonal, -1 on the $(i, i + 1)$ entries for $i = 1, \dots, d - 1$ and zeros elsewhere. If the matrix $\Lambda \in \mathcal{D}$, then the eigenvalues of the leading $(d - 1) \times (d - 1)$ submatrix of $\tilde{\Lambda}$ will all be negative.

For a set of d locations, one can supply the variogram matrix as valid input to the method.

9. **Brown–Resnick** (`br`) The Brown–Resnick process is the extension of the Hüsler–Reiss distribution, and is a max-stable process associated with the log-Gaussian distribution.

It is often in the spatial setting conditioned on a location (typically the origin). Users can provide a variogram function that takes distance as argument and is vectorized. If `vario` is provided, the model will simulate from an intrinsically stationary Gaussian process. The user can alternatively provide a covariance matrix `sigma` obtained by conditioning on a site, in which case simulations are from a stationary Gaussian process. See Engelke et al. (2015) or Dombry et al. (2016) for more information.

10. **Extremal Student** (`extstud`) of Nikoloulopoulos et al. (2009), eq. 2.8, with unit Fréchet margins is

$$P(\mathbf{X} \leq \mathbf{x}) = \exp \left[- \sum_{j=1}^d \frac{1}{x_j} T_{d-1, \nu+1, \mathbf{R}_{-j}} \left(\sqrt{\frac{\nu+1}{1-\rho_{ij}^2}} \left[\left(\frac{x_i}{x_j} \right)^{1/\nu} - \rho_{ij} \right], i \in I_{-j} \right) \right],$$

where T_{d-1} is the distribution function of the $d - 1$ dimensional Student- t distribution and the partial correlation matrix \mathbf{R}_{-j} has diagonal entry

$$r_{i,i;j} = 1, \quad r_{i,k;j} = \frac{\rho_{ik} - \rho_{ij}\rho_{kj}}{\sqrt{1-\rho_{ij}^2}\sqrt{1-\rho_{kj}^2}}$$

for $i \neq k, i, k \in I_{-j}$.

The user must provide a valid correlation matrix (the function checks for diagonal elements), which can be obtained from a variogram.

11. **Dirichlet mixture** (`dirmix`) proposed by Boldi and Davison (2007), see Dombry et al. (2016) for details on the mixture. The spectral density of the model is

$$h_{\mathbf{w}}(\mathbf{w}) = \sum_{k=1}^m \pi_k \frac{\Gamma(\alpha_{1k} + \dots + \alpha_{dk})}{\prod_{i=1}^d \Gamma(\alpha_{ik})} \left(1 - \sum_{i=1}^{d-1} w_i\right)^{\alpha_{dk}-1} \prod_{i=1}^{d-1} w_i^{\alpha_{ik}-1}$$

The argument `param` is thus a $d \times m$ matrix of coefficients, while the argument for the m -vector `weights` gives the relative contribution of each Dirichlet mixture component.

12. **Smith model** (`smith`), from the unpublished report of Smith (1990). It corresponds to a moving maximum process on a domain \mathbb{X} . The de Haan representation of the process is

$$Z(x) = \max_{i \in \mathbb{N}} \zeta_i h(x - \eta_i), \quad \eta_i \in \mathbb{X}$$

where $\{\zeta_i, \eta_i\}_{i \in \mathbb{N}}$ is a Poisson point process on $\mathbb{R}_+ \times \mathbb{X}$ with intensity measure $\zeta^{-2} d\zeta d\eta$ and h is the density of the multivariate Gaussian distribution. Other h could be used in principle, but are not implemented.

References

- Boldi, M.-O. (2009). A note on the representation of parametric models for multivariate extremes. *Extremes*, 12(3):211–218.
- Boldi, M.-O. and Davison, A. C. (2007). A mixture model for multivariate extremes. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(2):217–229.
- Coles, S. G. and Tawn, J. A. (1991). Modelling extreme multivariate events. *Journal of the Royal Statistical Society. Series B (Methodological)*, 53(2):377–392.
- Dombry, C., Engelke, S., and Oesting, M. (2016). Exact simulation of max-stable processes. *Biometrika*, 103(2):303–317.
- Engelke, S., Malinowski, A., Kabluchko, Z., and Schlather, M. (2015). Estimation of Hüsler–Reiss distributions and Brown–Resnick processes. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 77(1):239–265.
- Galambos, J. (1975). Order statistics of samples from multivariate distributions. *J. Amer. Statist. Assoc.*, 70(351, part 1):674–680.
- Gumbel, É. J. (1960). Distributions des valeurs extrêmes en plusieurs dimensions. *Publ. Inst. Statist. Univ. Paris*, 9:171–173.
- Huser, R. and Davison, A. C. (2013). Composite likelihood estimation for the Brown–Resnick process. *Biometrika*, 100(2):511–518.
- Hüsler, J. and Reiss, R.-D. (1989). Maxima of normal random vectors: between independence and complete dependence. *Statist. Probab. Lett.*, 7(4):283–286.
- Joe, H. (1990). Families of min-stable multivariate exponential and multivariate extreme value distributions. *Statistics & Probability Letters*, 9(1):75–81.
- Kotz, S. and Nadarajah, S. (2000). *Extreme value distributions*. Imperial College Press, London. Theory and applications.
- Nikoloulopoulos, A. K., Joe, H., and Li, H. (2009). Extreme value properties of multivariate t copulas. *Extremes*, 12(2):129–148.
- Pinheiro, J. C. and Bates, D. M. (1996). Unconstrained parametrizations for variance-covariance matrices. *Statistics and Computing*, 6(3):289–296.
- Smith, R. L. (1990). Max-stable processes and spatial extremes. Unpublished technical report.
- Stephenson, A. (2003). Simulating multivariate extreme value distributions of logistic type. *Extremes*, 6(1):49–59.
- Tawn, J. A. (1990). Modelling multivariate extreme value distributions. *Biometrika*, 77(2):245–253.